

**A PETITION TO THE STATE OF CALIFORNIA  
FISH AND GAME COMMISSION**

For action pursuant to Section 670.1, Title 14, California Code of Regulations (CCR) and Sections 2072 and 2073 of the Fish and Game Code relating to listing and delisting endangered and threatened species of plants and animals.

**I. SPECIES BEING PETITIONED:**

Common Name: Humboldt Marten

Scientific Name: *Martes caurina humboldtensis*

**II. RECOMMENDED ACTION:**

(Check appropriate categories)

a. List ☒

b. Change Status ☐

As Endangered ☒

from \_\_\_\_\_

As Threatened ☐

to \_\_\_\_\_

Or Delist ☐

**III. AUTHOR OF PETITION:**

Name: Environmental Protection Information Center, Center for Biological Diversity

Address: 145 G Street, Suite A, Arcata, CA 95521

Phone Number: (707) 822-7711

*I hereby certify that, to the best of my knowledge, all statements made in this petition are true and complete.*

Signature: Robert D. Pearson

Date: June 1, 2015

## **Before the California Fish and Game Commission**

**Petition to List the Humboldt Marten (*Martes caurina humboldtensis*) as an  
Endangered Species under the California Endangered Species Act**



**Environmental Protection Information Center and Center for Biological Diversity  
June 1, 2015**

# Table of Contents

I.	Executive Summary .....	3
II.	CESA Listing Process .....	3
III.	Population and Trends .....	4
IV.	Range and Distribution.....	6
V.	Abundance .....	8
VI.	Life History.....	9
VII.	Kind of Habitat Necessary For Survival .....	13
VIII.	Factors Affecting Ability to Survive and Reproduce.....	16
1.	Present or Threatened Modification or Destruction of Habitat.....	17
2.	Overexploitation .....	20
3.	Predation .....	22
4.	Competition .....	22
5.	Disease .....	23
6.	Other natural occurrences or human-related activities. ....	23
IX.	Degree and Immediacy of Threat.....	29
X.	Impact of Existing Management Efforts.....	30
XI.	Suggestions for Future Management.....	30
XII.	Availability and Sources of Information .....	31
XIII.	Detailed Distribution Map.....	38

## Notice of Petition

For action pursuant to Section 670.1, Title 14, California Code of Regulations (CCR) and Sections 2072 and 2073 of the Fish and Game Code relating to listing and delisting endangered and threatened species of plants and animals.

### I. SPECIES BEING PETITIONED

Common Name: Humboldt Marten

Scientific Name: *Martes caurina humboldtensis*

### II. RECOMMENDED ACTION: List as Endangered

The Environmental Protection Information Center and the Center for Biological Diversity submit this petition to list the Humboldt marten as an Endangered Species throughout its range in California pursuant to the California Endangered Species Act (California Fish and Game Code §§ 2050 et seq., "CESA"). This petition demonstrates that the Humboldt marten clearly warrants listing under CESA based on the factors specified in the statute.

### III. PETITIONERS

Primary Author: Rob DiPerna, California Forest and Wildlife Advocate, Environmental Protection Information Center

Address: 145 G Street, Suite A. Arcata, CA 95521

Phone Number: (707) 822-7711

E-mail: [rob@wildcalifornia.org](mailto:rob@wildcalifornia.org)

Additional Author and Petitioner: Center for Biological Diversity

Contact: Justin Augustine, 351 California St., Suite 600, San Francisco, CA 94104

Phone Number: 415-436-9682

E-mail: [jaugustine@biologicaldiversity.org](mailto:jaugustine@biologicaldiversity.org)

**The Environmental Protection Information Center** is a non-profit environmental advocacy organization with over 37 years of history protecting forests and species diversity in northwest California. [www.wildcalifornia.org](http://www.wildcalifornia.org).

**The Center for Biological Diversity** is a non-profit conservation organization with more than 775,000 members and online activists dedicated to the protection of endangered species and wild places. [www.biologicaldiversity.org](http://www.biologicaldiversity.org).

## I. Executive Summary

The Humboldt marten (*Martes caurina humboldtensis*) is a slender forest carnivore in the weasel family which is so rare that until recently it was thought to be extinct. Historically the Humboldt marten was abundant in coastal old-growth forests. This subspecies was once so common that it was regularly taken by trappers for its attractive fur. Due to historic trapping, drastic loss of old-growth forests from logging, and other stressors described below, the Humboldt marten has been extirpated from 95 percent of its historical range in California and from the vast majority of its range in Oregon. Only three populations remain. In northern coastal California, the sole remaining extant population likely totals around 40 individuals and is reproductively isolated from other coastal martens in Oregon and other subspecies of marten further inland. Oregon's two known populations, located in central and southern coastal Oregon, are likewise thought to be reproductively isolated and small in number. The isolated small populations compound the other stressors on the population, seriously endangering the Humboldt marten with extinction in California.

This petition summarizes the natural history of the Humboldt marten, its population status, and the ongoing threats to the subspecies and its habitat. Further, this petition demonstrates that, in the context of the CESA's six listing factors, the California Fish and Game Commission should list the Humboldt marten as endangered. Finally, this petition addresses existing management activities and recommends future management actions for the conservation of the Humboldt marten.

## II. CESA Listing Process

Recognizing that certain species of plants and animals have become extinct "as a consequence of man's activities, untempered by adequate concern for conservation," (Fish & G. Code § 2051 (a)), that other species are in danger of extinction, and that "[t]hese species of fish, wildlife, and plants are of ecological, educational, historical, recreational, esthetic, economic, and scientific value to the people of this state, and the conservation, protection, and enhancement of these species and their habitat is of statewide concern" (Fish & G. Code § 2051 (c)), the California Legislature enacted the California Endangered Species Act.

The purpose of CESA is to "conserve, protect, restore, and enhance any endangered species or any threatened species and its habitat." (Fish & G. Code § 2052). To this end, CESA provides for the listing of species as "threatened" and "endangered." The Commission is the administrative body that makes all final decisions as to which species shall be listed under CESA, while the CDFW is the expert agency that makes recommendations as to which species warrant listing.

For the purposes of CESA, a "threatened" species means "native species or subspecies of a bird, mammal, fish, amphibian, reptile, or plant that, although not presently threatened with extinction, is likely to become an endangered species in the foreseeable future in the absence of the special protection and management efforts required by this chapter. (Fish & G. Code § 2067). For the purposes of CESA, an "endangered" species means "a native species or subspecies of a bird, mammal, fish, amphibian, reptile, or plant which is in serious danger of becoming extinct throughout all, or a significant portion, of its range due to one or more causes, including loss of habitat, change in habitat, overexploitation, predation, competition, or disease." (Fish & G. Code § 2062).

The listing process may be set in motion in two ways: “any person” may petition the Commission to list a species or the CDFW may on its own initiative put forward a species for consideration. (Fish & G. Code § 2072.7). In the case of a citizen proposal, CESA sets forth a process for listing that contains several discrete steps. Upon receipt of a petition to list a species, a 90-day review period ensues during which the Commission refers the petition to CDFW, as the relevant expert agency, to prepare a detailed report. The CDFW’s report must determine whether the petition, along with other relevant information possessed or received by the Department, contains sufficient information indicating that listing may be warranted. (Fish & G. Code § 2073.5).

During this period interested persons are notified of the petition and public comments are accepted by the Commission. (Fish & G. Code § 2073.3). After receipt of CDFW’s report, the Commission considers the petition at a public hearing. (Fish & G. Code § 2074). At this time the Commission is charged with its first substantive decision: determining whether the petition, together with CDFW’s written report, and comments and testimony received, present sufficient information to indicate that listing of the species “may be warranted.” (Fish & G. Code § 2074.2).

This standard has been interpreted by courts as the amount of information sufficient to “lead a reasonable person to conclude there is a substantial possibility the requested listing could occur.” (*Natural Resources Defense Council v. California Fish and Game Comm.* (1994) 28 Cal.App.4<sup>th</sup> 1104, 1125). If the petition, together with CDFW’s report and comments received, indicates that listing “may be warranted,” then the Commission must accept the petition and designate the species as a “candidate species.” (Fish & G. Code § 2074.2).

### **III. Population and Trends**

Historically, Humboldt martens were common (Slauson et al. 2001). However, Humboldt martens in the coastal forests of Oregon and California have experienced significant declines, greater than those experienced by martens in the Sierra and Cascade mountains, because the narrow coastal forests are “more accessible to trappers, more accessible to logging, more rapidly affected by fragmentation of habitat and populations, and are composed of proportionately little late successional reserves or wilderness areas” (Slauson and Zielinski 2004, p. 63).

The Humboldt marten was thought to be either extremely rare or extinct (Kucera et al. 1995) until the late 1990’s, when martens were detected at two of 468 track plate stations within the range of the Humboldt marten by Zielinski and Golightly (1996). Prior to the redetection, Zielinski and Golightly (1996) were unable to verify a single marten detection within the historical range of the Humboldt marten since the 1940’s. Despite extensive survey efforts, martens have only been redetected in the north-central portion of their former California range and in central and southern coastal Oregon (Slauson and Zielinski 2007, Douglas and Holley 2009, Slauson et al. 2009b).

The Humboldt marten population is perilously small. The lone extant population of Humboldt marten in California appears to have declined by more than 40 percent from 2000–2008. Slauson et al. (2009a) report:

“The change between 2000–01 and 2008 marks a significant decline in site occupancy, equaling a change in occupancy rate ( $\lambda$ ) = 0.58 (SE = 0.13, 95% CI = 0.31 to 0.81) or a 42% decline in sample unit occupancy over the 7 year period.” (p. 10).

Because the researchers could not access all sites due to fires, the authors report that the decline in occupancy may likely be even higher than the reported 40 percent (Slauson et al. 2009a). Slauson et al. (2009a) estimated the population size in 2000–2001 at 31.5 individuals (95% C.I = 24–40), and in 2008 at 20.2 individuals (95% C.I = 11–30). Because they could not sample all potentially occupied habitat, this number could be an underestimate, and the authors estimate that even under the most optimistic scenario, the population in 2008 was likely somewhere around 40 individuals. Based on their survey results, Slauson et al. (2009a) concluded that immediate conservation actions are needed for the remnant California population. Based on subsequent surveys in 2012, Slauson et al. estimate that species abundance did not change in that time period. (USFWS 2015). In sum, based on the best available science, it is likely that there are less than 100 coastal Humboldt martens in California.

Less is known about the Oregon populations. According to the USFWS' 2015 Species Report:

[M]artens in coastal Oregon are currently known from two populations; one in central coastal Oregon and one in southern coastal Oregon. Slauson et al. stated that these two populations are small and isolated due to natural distribution of suitable habitat, historical and contemporary effects of timber harvest, and historical effects of fur trapping. Further, stated that marten populations in coastal Oregon and California have declined. Zielinski et al. reported that the number of martens harvested in coastal Oregon counties has declined since the 1940s. By the 1970s, martens were considered very rare along the Oregon coast. Abundance estimates for the two coastal Oregon populations are not available. Marten surveys in unsurveyed portions of coastal Oregon began in mid-2014, but the area surveyed to date represents a small amount of the currently available suitable habitat found within the CCO\_EPA and SCO\_EPA. Additional presence/absence surveys are required to accurately delineate the distribution of these two populations, and grid-based surveys are required to estimate abundance. Because U.S. Highway 101 runs the entire length of the range of martens in coastal Oregon and northern coastal California, scientists expect densities of road kills to reflect the abundance of martens. The best available data indicate an absence of reported marten road kills along coastal U.S. Highway 101 in northern coastal Oregon, southern coastal Oregon, and northern coastal California, suggesting low numbers of martens, at least within coastal habitat in the vicinity of U.S. Highway 101 in those regions.(Internal citations omitted).

The USFWS Species Report (2015) concludes, “[T]he fact that the two extant populations in coastal Oregon and single population in northern coastal California are considered to be small and isolated has led species experts (Slauson et al. 2009a, p. 1340) to have serious concerns about the viability of coastal marten populations.”

NatureServe (2015) ranks the Humboldt marten as an imperiled subspecies (T2S2) (last reviewed 2011). Kucera (1998) reports that *M. a. humboldtensis* “appears to meet CESA (California Endangered Species Act) criteria for listing as Endangered in its historic range of Del Norte, Humboldt, Mendocino, and Sonoma counties” due to historic trapping, habitat loss caused by logging, and severe population reduction (p. 141). Slauson et al. (2009b) report that they have serious concerns about the viability of coastal marten populations due to small population size, population isolation, and ongoing threats from logging.

## IV. Range and Distribution

For the purposes of listing under CESA, we will only discuss the range of the Humboldt marten in California. In California, the Humboldt marten historically occurred in coastal forests from Sonoma County, California, north to Curry County, Oregon (Slauson et al. 2001). Grinnel et al. 1937 described the range of the “well-marked race” as the coastal redwood (*Sequoia sempervirens*) zone from Sonoma County north to the Oregon border (Kucera 1998). There are natural heritage records for this subspecies from Colusa, Del Norte, Glenn, Humboldt, Lake, Mendocino, Siskiyou, Tehama, and Trinity counties in California (NatureServe 2015).

The Humboldt marten has been extirpated from more than 95 percent of its historic range in California (Slauson et al. 2007c). There is now only a single known population of this subspecies in the state (Zielinski et al. 2001, Slauson 2003, Slauson et al. 2007c). Slauson et al. (2007c) define the California study area for the Humboldt marten as portions of the Klamath-Siskiyou and Northern California Coastal Forest ecoregions in Del Norte, Humboldt, and Siskiyou counties, California. Slauson et al. (2009c) estimate that within its potential habitat, the single known California population of this subspecies currently occupies an area of 637 km<sup>2</sup>, using minimum convex polygon estimation. Based on a GIS analysis of Slauson’s data and more recent unpublished data, the Environmental Protection Information Center (EPIC) estimates the total California range of the subspecies to be approximately 2273 km<sup>2</sup> (Fig. 1).






Figure 1. Humboldt Marten California Range and Land Management. Map by EPIC GIS.

## V. Abundance

Table of distribution, abundance, and trend for the three extant population areas of the Humboldt marten in coastal Oregon and northern coastal California. Adapted from USFWS (2015).

Extant Population <sup>1</sup>	Total Area of Extant Population Area	Percent of Entire Historical Range	Abundance Estimate	Population Trend	Source(s) for Estimate
Central Coastal Oregon	4,150 km <sup>2</sup> (1,602 mi <sup>2</sup> )	7	No estimate	Presumed to be in decline <sup>2</sup>	Zielinski et al. (2001); Slauson and Zielinski (2009)
Southern Coastal Oregon	4,696 km <sup>2</sup> (1,813 mi <sup>2</sup> )	8	No estimate	Presumed to be in decline <sup>2</sup>	Zielinski et al. (2001); Slauson and Zielinski (2009)
Northern Coastal California	812 km <sup>2</sup> (313 mi <sup>2</sup> )	1.4	<100 individuals	Declining 2001–2008; unchanged 2008–2012 <sup>3</sup>	Slauson et al (2009a)

<sup>1</sup> All three marten populations in coastal California and Oregon are native. No known marten reintroductions have occurred in the analysis area.

<sup>2</sup> Both populations are believed to be in decline mainly on a reduction in the number of matens legally trapped and anecdotal sightings over time. However, current population numbers are unknown.

<sup>3</sup> The populations declined by approximately 42 percent between 2001 and 2008, based on a comparison of occupancy estimates from systematic grid-based surveys conducted in 2000–2001 and 2008. Preliminary occupancy estimates from 2012 are similar to 2008 values (Slauson, unpublished data).

## VI. Life History

### ***Species Description***

The marten is a forest-dwelling carnivore and a member of the weasel family (*Mustelidae*). Although martens are in the mustelid family with skunks and other species with powerful musk glands, the marten produces odors only weakly perceptible to humans (Buskirk and Ruggiero 1994). Martens have medium length glossy fur that ranges from tan to chocolate in color, an irregular cream or amber colored gular (throat) patch, lighter shaded fur on their underside, and darkly furred legs and tail. They are slender with a fox-like face, and large triangular-shaped ears that extend beyond the top of the head (Strickland et al. 1982). Of the two subspecies that occur in California, the Humboldt marten is darker with richer golden tone overall and with less orange and yellow in the throat patch (Grinnell and Dixon 1926, Buskirk and Zielinski 1999). It also has a smaller skull, and smaller and less crowded premolars and molars than the Sierra subspecies (Ibid.). One male Humboldt marten captured in mid-fall that had molted into winter pelage had brighter overall reddish brown coloration, dense fur on its underside, and dense fur around the pads of the feet (Slauson et al. 2002). Martens have five toes on each foot, all of which touch the ground, and semi-retractable claws (Buskirk 1994).

Adult American martens weigh from 500–1400 grams and are 500–680 mm long (Buskirk and McDonald 1989). Sexual dimorphism is pronounced, with males being 20–40 percent larger than females. Live capture data of 14 martens in the range of the Humboldt subspecies in coastal northwestern California show that martens in this geographic location are on the smaller end of the size range with an average weight of 889 grams for males (SD = 100), and an average weight of 598 grams for females (SD = 39) (Slauson et al. 2002).

### ***Taxonomy***

Marten taxonomy is in a state of flux. Petitioners believe that the best available science indicates that the Humboldt marten is a subspecies of *Martes caurina*.

The USFWS Species Report (2015) summarizes the current state of marten taxonomy:

A single species of marten, the American marten (*Martes americana*), was historically recognized as occurring across a broad range in North America, including the boreal forest region, montane coniferous forests, and Atlantic and Pacific coastal forested regions of North America. The Pacific marten (*M. caurina*) was recently split from the American marten based on genetic and morphological differences. The Pacific marten occurs largely in montane and coastal coniferous forests west of the Rocky mountain crest in North America, while the American marten occurs to the north and east of the Rocky mountain crest. The genetic split between these two species of martens is thought to have originated from the persistence of marten populations in two disjunct glacial refugia during the last glacial period.

In Oregon, two subspecies of martens have been historically recognized, with *M. caurina caurina* occurring in the Coast Range and Cascades Mountains of central and western Oregon and *M. c. vulpina* occurring in the Blue Mountains of northeastern Oregon. Two subspecies of Pacific marten occur in California.

The Humboldt marten (*M. c. humboldtensis*) occurs along the northern coast, whereas the Sierra marten (*M. c. sierrae*) is found in the interior mountains of northwestern California, the Cascade Mountains in northern central California, and the Sierra Nevada Mountains of eastern California.

The Humboldt marten was historically distributed throughout the coastal coniferous forests of northern California from northwestern Sonoma County northward to the Oregon border. Recent phylogenetic analyses using mitochondrial DNA (mtDNA) support the distinctiveness of the Humboldt marten subspecies, based on the presence of distinct haplotypes shared by historical museum specimens and martens currently occupying portions of the historical range in northern coastal California. Marten populations in coastal Oregon, which were historically described as *M. c. caurina*, also share these haplotypes, leading Slauson et al. (2009a) to suggest that martens in the Coast Range of Oregon may also be *M. c. humboldtensis*. Furthermore, preliminary results of a subspecific genetic evaluation of the Pacific marten by Schwartz et al. (In prep.)—using nuclear DNA (nDNA) and samples from substantially more martens than used by Slauson et al. (2009a)—also indicate that coastal Oregon and northern coastal California marten populations represent a single evolutionary clade calling into question the separation of the original subspecies range boundaries (i.e., *M. c. humboldtensis* in northern coastal California and *M. c. caurina* in coastal Oregon) at the California-Oregon border. (Internal citations omitted).

In sum, the best available science indicates that the coastal martens of California and Oregon are part of a discrete evolutionary clade. This genetic divergence, together with observed morphological differences between coastal California martens and other subspecies and presence in unique coastal forests, lends to the conclusion that the coastal martens of coastal California and Oregon are a subspecies of *M. caurina*.

### **Life History**

The life history traits of martens make recovery from population decline difficult. For a mammal of their size, martens have relatively late sexual maturity and low reproductive capacity. The predicted time to sexual maturity for a 1-kg mammal is five months (Taylor 1965), but most female martens first mate at 15 months of age and don't produce their first litters until 24 months (Strickland et al. 1982). During periods of environmental stress, pregnancy rates for marten can be as low as 50 percent, and females less than two years of age may not ovulate (Thompson and Colgan 1987). Martens produce only one litter per year, which is lower than the predicted frequency of 1.4 litters per year for similar sized mammals (Calder 1984). Litter size ranges from one to five with an average of 2.85 (Strickland and Douglas 1987), which is also low for a mammal its size (Calder 1984). Litter size is likely age-dependent, and peaks at age six. Senescence occurs at twelve years or greater, with decreased litter size beginning at twelve years (Mead 1994). Marten population densities are low, and are only about one-tenth the expected based on body size (Buskirk and Ruggiero 1994, Kucera 1998).

### **Reproduction**

Martens mate from late June to early August, with most mating occurring in July (Markley and Bassett 1942). Ovulation may be induced by copulation (Mead 1994). Like other species of Carnivora, marten undergo delayed implantation, and the active pregnancy period is only 27

days (Strickland et al. 1982). The onset of active gestation is controlled by photoperiod (Enders and Pearson 1943), and coincides with the development of mammarys (Mead 1994). Martens give birth in March and April, with newborn kits weighing approximately 28 grams (Strickland et al. 1982). Kits are weaned at 42 days, which is later than the predicted age for weaning of a 1-kg mammal of 28 to 34 days (Blaxter 1971, Millar 1977). At 50 days, young martens emerge from their dens and begin foraging independently (Hauptman 1979, Strickland et al. 1982). Juveniles disperse from early August to October. Martens are promiscuous, with both males and females having several mates, though it is unknown if multiple matings result in litters of multiple paternity (Strickland et al. 1982). Consistent with other polygynous Carnivora, only maternal care has been reported and includes establishing and maintaining the natal den, moving kits among alternative den sites, and grooming, nursing and bringing food to the young (Mead 1994). Martens produce an average of slightly less than three young per female with one litter per year (Strickland et al. 1982, p. 602). For a mammal of their size, martens have low reproductive rates and high longevity suggesting a slow recovery from population-level impacts (Buskirk and Ruggiero 1994, p. 16).

### **Longevity**

Martens can live up to 14.5 years in the wild (Strickland and Douglas 1987); however, as found in the USFWS Species Report (2015), “it appears that in the wild most North Americans marten, and presumably coastal martens, live less than 5 years.” There are numerous mortality factors for marten including predation, exposure, accidents, collision with automobiles, disease, and trapping. Bull and Heater (2001) reported that of 22 documented marten deaths in northeastern Oregon, 18 were killed by predators, 3 by exposure (hypothermia), and one in a collar-related accident. Of the 18 predator mortalities, 8 martens were killed by bobcats, 4 by raptors, 4 by other martens, and 2 by coyotes. Predation avoidance has likely influenced marten evolution and “led coastal martens to select for highly complex forest structure and avoid areas lacking overhead and escape cover.” (USFWS 2015). Mortality rates were approximately equal for males and females, though predation rates are generally higher for females, which are smaller (Slauson et al. 2009). Accidents may include falling out of trees and drowning (Buskirk and Ruggiero 1994).

### **Diet**

The marten is an opportunistic predator with a diverse diet that includes mammals, birds, carrion, eggs, insects, and vegetation (fruits, berries, nuts, fungi, lichens, grass, etc.) (Buskirk and Ruggiero 1994, p. 18; Martin 1994, p. 301). Voles (*Microtus* spp. and *Clethrionomys* spp.), squirrels (*Tamiasciurus* spp. and *Spermophilus* spp.), and chipmunks (*Tamias* spp.) are important food items for martens across their range (Martin 1994, p. 298). In the Sierra Nevada of California, mammals were the most important food item with microtine rodents the most frequent prey throughout the year and chipmunks and squirrels increasing in importance during the summer (Zielinski et al. 1983, p.388). Seasonal variation in diets is universal with the importance of soft mast, such as berries of *Vaccinium* and *Rubus* peaking in the fall (Buskirk and Ruggiero 1994, p. 19). Two key prey species in the winter diet, red-backed voles (*Clethrionomys californicus* and *C. gapperi*) and Douglas squirrel (*Tamiasciurus douglasii*), are closely associated with late-successional (mature and old-growth age classes) forest conditions (Slauson 2003, p. 6).

The USFWS 2015 Species Report notes:

Many of the key prey species of the Humboldt marten reach their highest densities in forest stands with old-growth structural features where their key

food resources—conifer seed crops and fruiting bodies of ectomycorrhizal fungi—reach their greatest abundances. The density of ericaceous (i.e., members of the plant family Ericaceae or heather family) shrub layers has also been shown to be positively correlated with chipmunk density in coastal Oregon. Physical complexity on or near the forest floor, typically provided by coarse woody debris, is directly related to predation success for martens; when this complexity is reduced (e.g., by logging), predation success declines due in part to the increased vigilance prey exhibit when physical complexity is reduced. (Internal citation omitted).

## ***Home Range***

The USFWS Species Report (2015) describes the Humboldt marten's home range:

Pacific and American martens exhibit strong habitat selection at the home range scale, suggesting that this scale of selection most directly influences an individual's fitness. Martens establish home ranges to encompass their year-round resource needs as well as, during the breeding season, access to members of the opposite sex. Theoretically, home range size for a predator is a function of prey density and habitat quality; smaller home ranges typically represent better habitat conditions. Marten home ranges are often positioned to maximize the composition of high quality habitat and minimize low quality habitat. Individual Pacific and American marten home ranges typically include a high proportion ( $\geq 70$  percent) of high quality late-successional forest habitat. Females, due to their solitary role raising young, have unique needs and must have access to reliable and nearby prey resources to support the energetic demands of lactation and providing food for kits. In northern coastal California, 97 percent (38 of 39) of a typical female's within-home range resting and active locations occurred in the core old-growth and late-mature riparian habitat patches. For males, 30 of 39 (77 percent) within-home range resting and active locations occurred in the core old-growth and late-mature riparian habitat patches.

There is an inverse relationship between the amount of high quality habitat and marten home range size. As the amount of low quality habitat (e.g., recent clear-cuts or partial harvests) increases, home range size increases. Accordingly, in our review of home range area studies of Pacific martens in California and Oregon, the largest home ranges we encountered (i.e., greater than 10 km<sup>2</sup> (3.9 mi<sup>2</sup>)) were from individuals occupying intensively logged landscapes.

Home ranges of Pacific martens in the Sierra Nevada Mountains of California in largely unlogged forest landscapes averaged 3–5 km<sup>2</sup> (1.2–1.9 mi<sup>2</sup>) for males and 3–4 km<sup>2</sup> (1.2–1.5 mi<sup>2</sup>) for females. Limited telemetry data from coastal martens suggests that home ranges for adult males ( $n = 3$ ) are of similar size (3–4 km<sup>2</sup>; 1.2–1.5 mi<sup>2</sup>). Telemetry data and habitat selection analysis at coastal marten detection sites reveal that home ranges include large patches (median  $> 1.5$  km<sup>2</sup> (0.6 mi<sup>2</sup>)) of the most favored habitat: old-growth, old-growth and late-mature, and serpentine habitat. (Internal citations omitted).

The USFWS Species Report (2015) provides the most up-to-date information on the home range size, selection, and dynamics of the Humboldt marten.

## VII. Type of Habitat Necessary for Survival

Martens are one of the most habitat-specific mammals in North America, and are thus highly vulnerable to habitat loss and degradation (Harris 1984, Buskirk and Ruggiero 1994, Slauson 2003). Martens are very strongly associated with closed-canopy, old growth forests with complex structure on or near the ground (Buskirk and Powell 1994, Buskirk and Ruggiero 1994, Bull et al. 2005). Martens are known to avoid younger forests and open areas such as clear-cuts (Drew 1995, Buskirk and Ruggiero 1994, Slauson et al. 2007). Martens avoid fragmented areas, and will not cross large areas with low canopy closure (Hargis and McCullough 1984, Bissonette and Sherburne 1993, Thompson and Harestad 1994, Hargis et al. 1999).

Numerous studies demonstrate the preference of martens for unlogged, old-growth habitat (Spencer et al. 1983, Wynne and Sherburne 1984, Snyder and Bissonette 1987, Koehler et al. 1990, Lofroth 1993, Buskirk and Ruggiero 1994, Raphael and Jones 1997, Ruggiero et al. 1998, Bull et al. 2005). For example, radio-collared martens in northeastern Oregon demonstrated a strong preference for old, unlogged stands with greater than 50 percent canopy closure, canopy layers, and high density of logs and dead trees (Bull et al. 2005). Martens specifically avoided harvested stands, early structural classes, and areas with low densities of dead trees (*Ibid.*). Younger forests and forests where old-growth loss has been extensive simply do not provide adequate habitat for martens—"The [Humboldt] marten does not occur in extensively logged redwood forests and currently only occurs in conifer-dominated, late-mature and old-growth forests with dense shrub cover or near-coast serpentine communities with dense shrub cover." (Slauson et al. 2003).

The Humboldt marten population documented in Slauson's 2003 study used two distinct types of fog-influenced coastal low-elevation forests—old-growth Douglas-fir (*Psuedotsuga menziesii*) dominated forests, and mixed-conifer forest occurring on serpentine soils with Douglas-fir, Sugar pine (*Pinus lambertiana*), Western white pine (*P. monticola*), and Lodgepole pine (*P. contorta*) (Slauson 2003). The more recent documentation of Humboldt marten presence in a third type of fog-influenced coastal low-elevation forest—old growth redwood forest—is consistent with both the earlier results and with the historical record (Grinnel et al. 1937, Kucera 1998, Slauson et al. 2001, Slauson and Holden 2009).

Serpentine soils have low levels of essential nutrients and high concentrations of detrital elements which creates a harsh growing environment and results in open and rocky sites with rich plant diversity, slow-growing woody plants, and stunted trees (Slauson et al. 2007). Serpentine areas provide lower quality habitat for martens than old-growth habitats (Slauson et al. 2009). In both serpentine and non-serpentine areas, the Humboldt marten occupies areas with dense, spatially extensive shrub cover comprised of shrub species associated with older forest habitats and importantly, not associated with shrub species that occur in areas of clearcuts and re-growth (Slauson et al. 2007). (*Ibid.*).

Martens select habitat at three spatial scales—microhabitat, stand, and home range, with a fourth scale, landscape, serving as an upper constraint on habitat selection (Bissonette et al. 1997, Slauson 2003). At all of these scales, martens demonstrate strong preference for old growth habitats.

## ***Microhabitat - Resting and Denning Structures***

The USFWS Species Report (2015)) describes microhabitat characteristics preferred by the Humboldt Marten:

Rest structures are used daily by martens between foraging bouts to provide thermoregulatory benefits and protection from predators. Reuse rates for individual rest structures are low and selection for structure type changes seasonally to meet thermoregulatory needs, such that multiple resting structures meeting seasonal requirements are required across the home range. Large-diameter live trees, snags, and logs provide the main types of resting structures for martens. Of 55 rest structures used by Humboldt martens in the summer and fall, 37 percent were snags, 23 percent downed logs, and 17 percent live trees. Martens typically select the largest available structures for resting and denning. Rest structures used by Humboldt martens averaged 95 cm (37 in.) diameter-at-breast-height (dbh) for snags, 88 cm (35 in.) maximum diameter for downed logs, and 94 cm (37 in.) dbh for live trees. These woody structures were found in the oldest forest development stages. Most resting locations—the actual place in the structure the marten used for resting—occurred in tree cavities (33 percent), on platforms (33 percent) created by broken top snags or large live branches, or in chambers (28 percent) created by log piles or rock outcrops. In coastal Oregon and northern coastal California, rest structures providing cavities or chambers will likely become seasonally important during the rainy period of the year; late fall through late spring.

Denning structures used by female martens to give birth to kits are called natal dens, and the subsequent locations where they move their kits are referred to as maternal dens. Pacific and American martens appear to be more selective of habitat conditions at den sites than at resting sites. Ruggiero et al. (1998) found that both the characteristics of the den structures and the characteristics of the stands they were found in influenced den-site selection. This is likely due to the importance of high quality foraging habitat in close proximity to den sites, allowing females to simultaneously maximize the energy they gain from foraging during lactation and minimize the time spent away from kits, especially when they are dependent on their mothers for thermoregulation. The most common den structures used by Pacific and American martens are large diameter live and dead trees with cavities. No natal dens and only three maternal dens (all from the northern coastal California population) have been described for the coastal marten. Two of the maternal dens were cavities in the broken tops of a 66 cm (26 in.) dbh golden chinquapin and 113 cm (44 in.) dbh Douglas-fir, and the other was a cavity in a 115 cm (45 in.) dbh Douglas-fir snag. All three maternal dens were located in the same old-growth, Douglas-fir dominated stand encompassing a creek and riparian habitat. (USFWS 2015). (Internal citations omitted).

The marten therefore requires structurally complex forest characteristics at the microhabitat level, especially snags and trees with cavities.



## ***Forest Stand Characteristics***

At the stand scale, which consists of several hectares, martens prefer old-growth stands with structural features that fulfill their life-history requirements such as resting and denning structures, abundant prey populations, access to mates, etc. (Buskirk and Powell 1994, Katnik et al. 1994, Slauson et al. 2007). Bull et al. (2005) compared habitat characteristics in 2,558 plots in occupied and unoccupied areas in northeastern Oregon and found that marten use stands with 50–74 percent canopy closure more than stands with less than 50 percent canopy closure ( $P < 0.01$ ), and that stands used by martens had more canopy layers, a longer distance to an opening, and higher densities of snags, logs, and large trees than unused areas. Importantly, they found that stands with no harvesting activity were used more, and that stands with *any* harvesting activity were used less than expected based on availability ( $P < 0.01$ ; emphasis added).

In non-serpentine stands, Humboldt martens used late-successional stands highly disproportionate to their availability, used late-mature stands similar to availability, and made little or no use of all other seral stages (Slauson et al. 2007, p. 462). All earlier seral stages were selected against, probably because of the lack of one or more key structural features (Slauson 2003, p. 62).

## ***Home Range Scale***

Humboldt martens select the largest available patches of old-growth, old-growth and late-mature, or serpentine habitat (Slauson et al. 2007), similar to home-range selection for other marten subspecies which also select for the largest available forest patches (Chapin et al. 1998). Slauson et al. (2007) developed habitat models for Humboldt marten and found that a 20-ha increase in old-growth patch size was associated with a 19-26 percent increase in marten occurrence, after accounting for the amount of serpentine habitat. They conclude, “The best models suggest that home range areas with larger patch sizes of old-growth, old-growth plus late-mature, or serpentine habitat within a 1-km radius of each sample unit are important for marten occurrence. Martens disproportionately used sample units within these largest patch sizes.”

Slauson et al. (2007) also found that mixed-scale models from both the stand and home range scales best explained Humboldt marten occurrence compared to one scale alone. Because martens are negatively associated with logging activities at the microhabitat, stand, and home-range scale, it follows that logging at the landscape scale, which is comprised of tens to hundreds of square kilometers, inevitably negatively influences marten occurrence.

## ***Landscape Scale***

Loss and fragmentation of mature forest at the landscape scale and the resultant changes in landscape pattern constrain marten movement and demography (Bissonette et al. 1989, Fredrickson 1990, Phillips 1994, Chapin 1995, Chapin et al. 1998, Hargis 1996, Slauson 2003). Martens avoid landscapes where more than 25–30 percent of mature forest has been lost (Bissonette et al. 1997, Hargis et al. 1999, Potvin et al. 1999, Slauson 2003). Fragmented forests and small patches of old growth are not adequate to ensure the long-term viability of marten populations. Slauson et al. (2009a) found that Humboldt martens may occasionally occupy patches of old growth forest that are less than 50 ha, but that these patches do not provide the same value to martens as larger patches: “The biggest difference between sites with stable marten occupancy versus unstable occupancy, in our study, was

the size of the patch of Old Growth forest that encompassed them, with larger patches having more stable marten occupancy.”

Slauson et al. (2009a) documented a sharp decline in Humboldt marten sample unit occupancy in sample units where the old-growth vegetation type is highly fragmented and in serpentine areas, and found that sample units with more old growth in their vicinity were more likely to have stable marten occupancy between 2000–01 and 2008 (Slauson et al. 2009a). Old growth patches encompassing sample units where marten occupancy remained constant between 2000-01 and 2008 were approximately 40 percent larger on average than patches that became unoccupied in 2008 (Slauson et al. 2009a). The authors calculated the probability of extinction for the marten based on old-growth patch size and found that martens were less likely to go extinct in sample units with more old growth. For example, a 30 hectare increase in the amount of old growth resulted in a 37 percent decrease in the probability of extinction.

As summarized by the USFWS (2015):

Compared to other species closely associated with late-successional forest, American and Pacific marten populations, including the coastal marten, are very sensitive to the loss or fragmentation of high quality habitat at the landscape scale. Martens exhibit a progression of responses to timber harvest as the proportion of habitat affected by intensive logging practices increases, including (but not limited to) activities such as clear cutting, partial harvest, and shelterwood cutting. The combination of habitat loss and fragmentation of remnant suitable habitat effectively lowers the density of martens by reducing the number of home ranges that can be supported. (USFWS 2015) (Internal citations omitted).

Patch size and degree of connectivity have important implications for martens attempting to recolonize unoccupied stands. Highly fragmented forests may contain suitable habitat patches that are so separated by open areas that martens cannot make use of the habitat that is available (Buskirk and Powell 1994, p. 289). The more highly fragmented mature forest becomes, the lower the carrying capacity for martens (Thompson and Harestad 1994, p. 360).

## **VIII. Factors Affecting Ability to Survive and Reproduce**

The Humboldt marten warrants listing under CESA. As described in this section, the Humboldt marten is threatened by all six factors identified by CESA as indicators that listing is warranted. The six listing factors to be addressed, as identified by CESA include:

- (1) Present or threatened modification or destruction of its habitat;
- (2) Overexploitation;
- (3) Predation;
- (4) Competition;
- (5) Disease; or
- (6) Other natural occurrences or human-related activities.

## 1. Present or Threatened Modification or Destruction of Habitat

### ***Timber Harvest and Logging***

As noted elsewhere in this petition, the Humboldt marten has been extirpated from the vast majority of its historic range. The primary cause of marten population decline and extirpation is loss of old-growth coniferous forests (Yeager 1950, Archibald and Jessup 1984, Thompson and Harestad 1994, Bull et al. 2005). Logging, by its very nature, necessarily threatens martens and their habitat because martens select the largest and oldest trees available at the microhabitat, stand, home range, and landscape scale (Wilbert 1992, Buskirk and Ruggiero 1994, Gilbert et al. 1997, Raphael and Jones 1997, Ruggiero et al. 1998, Bull et al. 2005, Slauson et al. 2007, 2009).

The USFWS Species Assessment (USFWS 2015) discusses the impacts of timber harvest activities on marten habitat:

Historically, in California, a primary reason for the range reduction of the coastal marten was likely the result of habitat loss due to logging of late-successional forests during the last century. Redwood accounted for approximately 35 percent of the conifer forests in the historical range in California. Zielinski et al. concluded that the effect of past and current timber harvest in the redwood region is the most plausible reason for the continued absence of the coastal marten throughout most of northern coastal California. Similar to northern coastal California, much of coastal Oregon is privately owned and the majority of late seral coastal forest stands have been harvested over the past century, especially in stands nearest to the coast. Little or no suitable marten habitat occurs in these privately owned areas adjacent to the coast. Most currently suitable coastal marten habitat near the coast in Oregon and California is federally owned. (USFWS 2015) (Internal citations omitted).

The majority of coastal forests on private lands have been logged at least once. Private forests in the range of the Humboldt marten are logged primarily by clear-cutting, and are currently managed under short rotation even-aged silvicultural regimes (60-70 years) which create structurally simplified, early to mid-seral landscapes that do not support martens (USDA 1992, Bolsinger and Waddell 1993, Lettman and Campbell 1997, Thornburg et al. 2000).

The USFWS Species Assessment (USFWS 2010) describes the importance of late-successional forests and forest structures for the marten:

Humboldt martens use structural features of late-successional forests, such as large diameter live trees, snags, and logs. Most resting structures used by Humboldt martens require more than a century to develop. Loss of these elements can reduce the suitability of forested areas for martens. The probability of detecting Humboldt martens increased with increasing maximum patch size of late-successional forest. The minimum patch size to identify potential Humboldt marten home range areas is 445 ac (180 ha) of late-successional forest with dense shrub cover. Little habitat with the necessary structural characteristics for Humboldt martens is expected to regenerate over the next several decades. (USFWS 2010, p. 17) (Internal citations omitted).

The USFWS 2010 Species Assessment Report further describes the negative effects of timber harvest on the marten and marten habitat:

[R]eduction in the total amount of late-successional forest is only one of the timber harvest-related threats facing the Humboldt marten. The continued simplification of the structure of forests and their fragmentation into smaller, more isolated, patches is also a concern. For example, RNSP contain approximately 41,400 ac (16,700 ha) of late-successional coast redwood forest. This includes 41,100 ac (16,600 ha) in stands greater than 0.5 ac (0.2 ha) and 362 ac (146 ha) of isolated late-successional trees surrounded by mature tree buffers. Late-successional forests in RNSP are fragmented with only three stands containing more than 5,000 ac (2,023 ha). The majority (83%) of the late-successional stands in RNSP are 100 ac (40 ha) or smaller with 31% of the stands less than 5 ac (2 ha) in size. RNSP also contains over 50,000 ac (20,235 ha) of second growth forest most of which was harvested between 1950 and 1978. (USFWS 2010, p. 17) (Internal citations omitted).

Federal lands in the range of the Humboldt marten have been managed under the Northwest Forest Plan since 1994. NWFP land allocations within the occupied Humboldt marten area include matrix, wilderness, Late Successional Reserves (LSRs), and administratively withdrawn areas. Matrix lands and LSRs are available for future timber harvest, while wilderness and administratively withdrawn areas are generally not. Approximately 38% of the occupied range occurs on public or private lands currently available for timber harvest.

The USFWS 2010 Species Assessment describes the challenges with relying on public lands to conserve the Humboldt marten:

Public lands are disjunct and represent only a small proportion of the total area of coastal forest in northern California, with most forestland in industrial or non-industrial private ownership. Public lands in coastal northwestern California include state parks, one national park, a Bureau of Land Management conservation area, and portions of the Six Rivers and Klamath National Forests. Humboldt martens were detected proportionately more frequently on lands managed by the Forest Service than on private timberlands. Greater than 80% of the private timberlands in the 2000- 2001 Humboldt marten study area were logged prior to surveying and martens appeared to avoid all but the edge of this landscape (USFWS 2010, p.17) (Internal citations omitted).

Sufficient evidence exists to indicate that the petitioned action 'may be warranted' in light of the threats posed by timber harvest activities on federal and non-federal lands. Timber harvest and logging are a present and ongoing threat to the survival and conservation of the Humboldt marten, especially in light of the marten's extremely small population size in California and the isolation of this population. Existing forest management regimes fail to take into account the needs of the Humboldt marten. (See also *Inadequacy of Regulatory Mechanisms, and Impacts of Existing Management Efforts*).

### ***Impacts of Fire, Fire Suppression, and Post-fire Salvage Logging on Marten Habitat***

Over the long-term, wildfire plays a role in developing the habitat components on which martens depend, such as snags and dense shrubs, but because the marten's habitat has been so severely reduced due to logging, fires can pose a threat to the subspecies. Due to its

critically low population size and restricted range, fire can potentially result in short-term loss and fragmentation of suitable habitat (Slauson and Zielinski 2004). Slauson (2003) identifies the risk of being extirpated by a stochastic event as a “major challenge” to the persistence and restoration of the Humboldt marten (p. 71). Commercial timber harvest (including salvage logging) and fire suppression have exacerbated the threat posed to the marten by fire by fragmenting landscapes.

The USFWS 2010 Species Assessment also describes the potential hazards to the marten resulting from fire suppression activities:

[T]he Humboldt marten is also threatened by fire suppression activities. Fuels management projects designed to lower fire risks, such as mechanical treatments have the potential to negatively affect Humboldt martens and their habitat by reducing important features such as shrubs, canopy cover, snags, or logs. Significant loss of the shrub layer due to fuels reduction projects may reduce habitat suitability, due to reduction in prey abundance or improved access by competitors. (USFWS 2010) (Internal citations omitted).

Post-fire salvage logging and hazard tree removal occurs on both federal and non-federal lands within the range of the Humboldt marten and can fragment marten habitat and eliminate important structure such as snags and large dead trees. While the impact of post-fire logging on the Humboldt marten has not yet been examined, recent science on the closely-related Pacific fisher shows that the fisher can utilize and may *prefer* higher-severity post-fire areas. (Hanson 2013, 2015). While a fire may remove shrub and canopy cover in the short term, post-fire forests contain high snag, shrub, and downed large woody debris density, providing a complex structural environment. This “complex early seral forest” is “rich in native biodiversity and wildlife abundance,” including small mammals, a staple of the marten’s diet. (Hanson 2013, 2015). Post-fire logging removes structural complexity, degrading the habitat potential of post-fire forests.

On federal lands, salvage logging and other fuels management activities can occur in all land allocations with the exception of wilderness areas. (USFWS 2010, p. 20). On non-federal lands, salvage logging and hazard tree removal activities are primarily conducted pursuant to emergency exemption notifications. While emergency exemption notices must be filed with the California Department of Forestry and Fire Protection (“CAL FIRE”), approval of such emergency timber operations notifications is not subject to the normal provisions of the California Environmental Quality Act (“CEQA”) review process. Rather, emergency exemption notification approval is considered to be a ministerial action, thus leaving CAL FIRE with little to no discretion. The emergency exemption process also excludes the possibility of public review and comment. (*See also Inadequacy of Regulatory Mechanisms, and Impacts of Existing Management Activities*).

In summary, the Humboldt marten is threatened by past, ongoing, and proposed habitat loss and modification due to logging and timber harvest, wildfire, fire suppression activities, and post-fire salvage logging. The existence and extent of these threats clearly demonstrates that listing of the Humboldt marten under CESA is warranted.

### ***Collisions with Vehicles***

Vehicular collisions are also a significant threat to the Humboldt marten, particularly given the small and isolated populations. As stated in the USFWS Species Report (2015):

Collision with vehicles is a known source of mortality for the coastal marten currently and [is] expected to continue into the future given the presence of roads within the range of the species. Collisions with vehicles may negatively affect the viability of the three coastal marten populations if annual roadkill mortality rates, in combination with all other sources of mortality, exceed annual juvenile recruitment rates. Based on their small body size, we expect that nearly all coastal martens struck by vehicles will either die immediately from blunt force trauma, or soon thereafter from severe injuries. A small proportion of coastal martens may survive a vehicle strike, but would likely be physically impaired, potentially increasing the probability of mortality from another source such as predation, disease, or starvation. For example, a broken limb from a vehicle strike would reduce the foraging capability of a coastal marten, which relies on swift movements to catch highly mobile prey through rugged terrain and into trees. If the injured coastal marten is an adult female that is caring for dependent kits, her reduced ability to acquire adequate prey may lead to one or more of the kits starving.

The lack of highway structures to facilitate free movement and connectivity of habitats for the Humboldt marten is a significant barrier to marten survival and dispersal. Vehicular collisions pose a significant threat to the Humboldt marten.

## **2. Overexploitation**

As found in the USFWS Species Report (2015):

Extensive unregulated fur trapping conducted over a long period of time can lead to a species range reduction, the extirpation of populations, or extinction of the species. Unregulated coastal marten fur trapping beginning in the late 1800s led to a marked reduction in the species distribution across coastal Oregon and northern coastal California by the late 1920s (see details below). Localized trapping may also negatively affect populations through removal of individual coastal martens that are vital to the long-term viability of the population (e.g., reproductive adult females). Population level effects of coastal marten trapping have not been studied, but coastal marten population growth is most significantly affected by a reduction in marten survival rates . . . Therefore, for small coastal marten populations, the loss of only a few adult martens each trapping season could reduce the likelihood of long-term population viability into the future, especially when combined with mortalities from other sources such as roadkill, disease, predation, and exposure to toxicants. Annual juvenile recruitment (estimated at 50 percent for the coastal marten; Slauson et al. In prep. (a)) may offset losses due to legal trapping. However, annual mortality rates from other sources are either unknown (e.g., roadkill, disease, exposure to toxicants) or derived from a relatively small area and small sample size of martens.

By the late 1800s and early 1900s, European settlers began trapping for the fur trade industry. By the early 1900s, annual harvest totals of martens in the analysis area were already in decline, signaling stress on populations from trapping. Accounts of individual trappers taking 35 and 50 martens in single winters within the California portion of the analysis area indicate the impact individual trappers had on marten populations. The sharp decline in annual harvest rates prompted Dixon to call for the closing of the marten trapping

season in California or fear of their extirpation. However, marten trapping continued and further reduced populations, resulting in excessive harvest of coastal marten populations that will take many years to recover and likely resulting in the loss of genetic variation .

The number of martens harvested in coastal Oregon counties has declined since the 1940s, and by the 1970s martens were considered very rare along the Oregon coast. Historical trapping of coastal martens for fur is considered by researchers as the likely cause of the marked contraction in coastal marten distribution and reduction in population size that was observed in coastal Oregon and northern coastal California in the early 20th century. The trapping season for martens was closed in 1946 in the California portion of the analysis area; however, decades of protection from trapping have not resulted in the recovery of coastal marten populations in northern coastal California.

Currently, trapping for martens is illegal in California. However, it is legal to trap other mammals that may occur within occupied coastal marten habitat in northern coastal California (e.g., bobcat, gray fox). . . .

In contrast to California, trapping of coastal marten for their fur is currently legal in Oregon. Coastal marten trapping records for the Oregon portion of the analysis area peaked in the 1940s and in no decade since have coastal marten harvest levels reached  $\geq 15$  percent of the 1940s total coastal marten harvest. Currently, the harvest of marten in the Oregon Coast range is rare. For example, three coastal marten were trapped within the entire Oregon portion of the analysis area (and area of 36,348-km<sup>2</sup> (14,034 mi<sup>2</sup>)) during the 2013 trapping season. A total of 36 martens (mean = 2.7 harvested per year; range = 0–5; standard deviation = 1.90) were harvested within coastal Oregon counties between 1969 and 1995. We excluded martens harvested from 1969 to 1995 in Lane (310 martens harvested) and Douglas counties (167 harvested) since both of these counties extend from the coast to the Oregon Cascades. Although most martens harvested in those two counties were likely from the Oregon Cascades, the actual proportion harvested within the Coast Range is unknown, but expected to be similar to the small number harvested in other coastal counties during that time frame.

The low number of coastal marten trapped in the Oregon portion of the analysis area in the recent past could be due to low marten densities in the two extant population areas in coastal Oregon, or possibly due to low trapping effort. Few Oregon trappers (4 to 8) have pursued marten in recent years, with most marten captures occurring in the Cascade Range of interior Oregon, which is outside the historical range of the coastal marten. The three coastal martens (two males and one female) harvested during the 2013 trapping season were all captured within 2 km (1.2 mi.) of the coast between 8 and 17 km (5 and 11 mi) north of Coos Bay, Coos County, Oregon). The proximity of all three harvested coastal marten to one of the more heavily [human] populated areas (i.e., Coos Bay, Oregon) in the [central coastal Oregon extant population area] suggests that trapping activity might be more prevalent near human population areas that are readily accessible by vehicles, such as via U.S. Highway 101 that traverses the [central coastal

Oregon extant population area] north to south near the coastline. (USFWS 2015). (Internal citations omitted).

While the threat posed to the Humboldt marten by accidental capture and poaching in California may be small, any loss to the marten population in California is significant due to the small population size and population isolation of the marten in California. Incidental trapping of Humboldt martens remains a threat to the survival and conservation of the species, and is yet another factor which demonstrates that listing under CESA is warranted.

### **3. Predation**

Predation is a significant threat to the Humboldt marten. Predation is a primary source of marten mortality. Bull and Heater (2001) report that of 22 documented marten deaths in their study in northeastern Oregon, 18 of the martens were killed by predators (82 percent). Martens face many predators including bobcats, foxes, coyotes, mountain lions, great horned owls, goshawks, and Pacific fishers. (Buskirk and Ruggiero 1994, Bull and Heater 2001, Slauson et al. 2009).

Habitat degradation and fragmentation caused by logging increases the threat of predation for martens by favoring generalist predators which fare better in logged landscapes (Slauson et al. 2009). As found by Slauson and Zielinski (2007), the distribution of mesocarnivores in redwood forests has changed over the last 80 years; fisher and gray foxes have expanded their distributions into Humboldt marten habitat concurrent with the dramatic decline of the marten. Slauson and Zielinski (2010) have further found that roads may be facilitating the increased presence of mesocarnivores in dense-shrub landscapes that martens prefer.

Higher predation rates attributable to habitat fragmentation and degradation may suppress marten populations. Slauson et al. (2009) found a higher decline in Humboldt marten sample unit occupancy from 2001–2008 in serpentine habitats and in sample units where old-growth is more fragmented and the risk of predation is increased.

Slauson et al. (2009) also found that sample unit occupancy declined more dramatically in units occupied by only female martens than in units with dual-gender or male-only occupancy. Because body size of female martens is generally 40 percent smaller than males, females are likely more vulnerable to predation from larger-bodied mesocarnivores associated with early seral and fragmented landscapes (*Ibid.*). Due to the extensive loss and fragmentation of old-growth forest habitats in coastal forests in California and Oregon and the resulting habitat conditions which favor marten predators, the Humboldt marten faces heightened predation threat. (*Ibid.*).

In summary, while predation is a natural stressor, human activities such as vegetation management contribute to this stressor and amplify it, and it therefore represents a significant threat to the Humboldt marten.

### **4. Competition**

The USFWS 2010 Species Assessment identifies environmental fluctuations due to variations in predation, disease, and food supply as a threat to the Humboldt marten as a consequence of its small population size in California. (USFWS 2010, p. 22). No data or studies have been produced to assess the impacts of inter-species competition on the Humboldt marten. However, given the precariously small population size, the threat of ongoing and habitat loss and fragmentation that results in favorable conditions for generalist predators, and the effects of climate conditions and drought on availability of preferred prey



species, competition for space and food with other mesocarnivores and other predators, competition is likely a currently a limiting factor for the ability of the Humboldt marten to survive and reproduce. The potentially detrimental effects of inter-species competition is yet another reason demonstrating that listing under CESA is warranted.

## **5. Disease**

The threat posed to the Humboldt marten by disease has not been studied, but given the subspecies' small population size, is potentially critical. As concluded by the USFWS, "The outbreak of a lethal pathogen within one of the three coastal marten populations could result in a rapid reduction in population size and distribution, likely resulting in a reduced probability of population persistence, given the small size of these populations." (2015).

Martens are susceptible to several mesocarnivore diseases and parasites including but not limited to rabies, plague, distemper, toxoplasmosis, leptospirosis, trichinosis, sarcoptic mange, canine adenovirus, parvovirus, and herpes virus, West Nile virus, and Aleutian disease (Strickland et al. 1982, Banci 1989, Green et al. 2008, IERC 2008). Though little information on the threat posed to the Humboldt marten by disease is available, many of the diseases to which it is susceptible are known to cause mortality in carnivores (e.g. Brown et al. 2008). Brown et al. investigated pathogen exposure in the Pacific fisher, a closely-related species, on the Hoopa Valley Indian Reservation. The Reservation is located near occupied marten habitat, only 9km (5.6 mi) south. Of the 15 fishers found dead on the Reservation, two had been exposed to the canine distemper virus and six to canine parvovirus. Secondary infections resulting from intra-species fighting or encounters with other larger predators may also cause mortality.

In summary, disease has the potential to be a threat due to the extremely small population size of the marten. While further study of susceptibility of the Humboldt marten to disease is required, it is clear that the threat of adverse impacts of disease to such a small population is a factor that demonstrates that listing under CESA is warranted.

## **6. Other natural occurrences or human-related activities.**

The Humboldt marten is threatened by several additional natural occurrences and human-related activities. These include: (a) Inadequacy of Regulatory Mechanisms (b) Toxic Exposure; and (c) Climate Change. This petition addresses each of these additional factors which demonstrate that listing of the Humboldt marten under CESA is warranted.

### **(a) Inadequacy of Regulatory Mechanisms**

Existing regulatory mechanisms are not adequate to protect and restore the habitat the marten needs to survive. Much marten habitat is currently available for logging, and further loss or degradation of its limited suitable habitat could push the Humboldt marten to extinction. Marten conservation will require landscape management to enlarge and reconnect suitable patches of habitat (Slauson et al. 2007). Conservation measures that aim only to maintain current marten habitat conditions will not ensure the Humboldt marten's long-term persistence (Slauson 2003).

There are no laws or regulations in California that adequately address the needs of the Humboldt marten on state, federal, non-federal, or tribal lands. Though the marten is protected from fur trapping in California, there are currently no regulatory mechanisms that adequately protect habitat for the Humboldt marten in the state.

The vast majority of coastal old-growth forests in California have been logged, causing drastic loss and fragmentation of marten habitat (Slauson 2003). Logging negatively affects martens at the microhabitat, stand, home range, and landscape scale. (*See also Kind of Habitat Necessary for Survival*). To this day, there are no laws or regulations that prohibit marten habitat loss and fragmentation due to logging of old growth or late successional forest stands on federal, non-federal or tribal lands.

### ***Federal Lands***

The Humboldt marten occurs on federal lands managed by the U.S. Forest Service and the National Park Service. The Forest Service manages the majority of the marten's range (Slauson et al. 2007, EPIC 2010). The range of the Humboldt marten includes the Six Rivers National Forest and Klamath National Forest in California, the Siskiyou National Forest in southwestern Oregon, and the Siuslaw National Forest in the central Oregon Coast Range. In California (Region 5), the marten is a Forest Service Sensitive Species, and was recognized as a priority species in Fiscal Year 2007.

Sensitive Species status, even as a priority species, does not afford the marten or its habitat the protection it needs to survive. Sensitive Species are not necessarily afforded any regulatory habitat protection; rather the agency is only required to analyze the impacts of its actions on the marten under the National Environmental Policy Act (NEPA). This requirement in no way mandates the agency to select an environmentally benign alternative or to try to mitigate the adverse impacts of projects. Moreover, any protections afforded the marten under the Sensitive Species program are discretionary. Discretionary mechanisms are not adequate to protect the marten on National Forest lands because National Forests are managed to meet multiple objects including providing access to recreation opportunities for the public and serving as an economic development resource for the regions where they occur (e.g. Six Rivers National Forest 2010).

Much of the marten's range on National Forests is managed under the Northwest Forest Plan (USDA/USDI 1994a, 1994b). The NWFP created seven types of land allocations including Congressionally Reserved Areas, Late-Successional Reserves, Managed Late-Successional Areas, Adaptive Management Areas, Administrative Withdrawn Areas, Riparian Reserves, and Matrix lands, each with different management guidelines. Though matrix lands harbor some of the remaining old growth forest in the range of the marten, these lands were intended to provide for commercial timber harvest rather than to provide wildlife values. Slauson (2003) detected martens on 8 of 31 sample units in matrix lands, where the potential for timber harvest poses a dire threat to its persistence. Of the land managed by the Forest Service in the range of the Humboldt marten surveyed by Slauson et al. (2007), 20 percent was designated as matrix land that is currently available for logging. Sixteen percent of the matrix land has already been logged (Slauson et al. 2007).

Under the NWFP, LSRs were intended to support viable populations of late successional and old growth associated species, but some rare species, including the marten, are not effectively protected by the reserve system. Moreover, logging is not prohibited on LSRs and therefore the potential for logging on these lands is a further threat to the marten. Of the FS land in the range of the Humboldt marten surveyed by Slauson et al. (2007), 40 percent was designated as Late Successional Reserve. Late Successional Reserves are not necessarily in late-seral condition, but are in theory being managed to develop mature forest conditions over time. These reserves do not necessarily provide habitat benefit to the marten currently because the conditions which marten prefer may take centuries to develop. Slauson (2003)

detected martens at 13 of 66 sample units in Late Successional Reserves. Thirteen percent of the Late Successional Reserves in the marten's range have been logged (Slauson et al. 2007).

Though the status of the marten was considered during the planning process for the NWFP, the process "did not include significant review of existing data or collection of new data" concerning the marten (Zielinski et al. 2001). The marten was given the second poorest score among mammals by the Forest Ecosystem Management Scientific Analysis Team for likelihood of remaining well distributed, with only a 67 percent likelihood of remaining well distributed within the range of the northern spotted owl (USDA/USDI 1994). Even this bleak projection, however, was overly optimistic, and Slauson et al. (2009a) conclude that the Northwest Forest Plan has not proven adequate to protect the Humboldt marten:

In reality, the situation is far worse, martens on federal lands in the Coast Range of California are restricted to a single refugia and have been extirpated from a significant portion (>95%) of their historical range. Within their last stronghold, measures including the protection of Riparian Reserves, Late-Successional Reserves, northern spotted owl and marbled murrelet conservation measures, do not completely protect the population (p. 3).  
(*Ibid.*).

Slauson et al. (2009a) report that at least 38 percent of the distribution of martens in coastal California occurs outside of NWFP reserves, and the reserves themselves may not contain suitable old growth habitat, as discussed above.

The Forest Service also manages the Siskiyou Wilderness, which is administered by the Six Rivers, Klamath and Siskiyou National Forests. The proportion of the marten's range which is designated as wilderness is not sufficient in and of itself to provide enough habitat to ensure long-term marten persistence for several reasons.

First, designated wilderness makes up only a small portion of the marten's range. EPIC (2010) estimates that 14 percent of the California range of the Humboldt marten consists of designated wilderness. Of the FS land in the range of the Humboldt marten surveyed by Slauson et al. (2007), 18 percent was designated as wilderness.

Second, not all vegetation types in the Siskiyou Wilderness support martens. Slauson (2003) detected martens at only 3 of 23 sample units in wilderness. Much of the Siskiyou Wilderness is composed of higher elevation vegetation such as white-fir and hardwood-dominated stands which are not preferred by the Humboldt marten (Slauson 2003).

In addition, the Forest Service manages the Smith River National Recreation Area, which is part of the Six Rivers National Forest. Occurrence in the NRA is not adequate to protect the Humboldt marten because management of National Recreation Areas prioritizes recreational opportunities over wildlife values. Though the habitat of the marten in the NRA is not vulnerable to timber harvest, it remains vulnerable to other threats. EPIC (2010) estimates that the Smith River NRA makes up 9 percent of the California range of the Humboldt marten.

The Humboldt marten also occurs on federal lands managed by the National Park Service. The Redwood National Parks complex consists of a series of parks managed by the National Parks Service and California State Parks including Prairie Creek, Jedediah Smith, and Del Norte Coast Redwoods State Parks. Martens were not known to be extant in the parks until

2009 when a marten was detected in Prairie Creek Redwoods State Park via a remote sensing camera (Slauson and Holden 2009). Habitat in the parks is not sufficient to ensure the survival of the Humboldt marten because the parks make up only a small portion of the marten's range, the parks do not currently support a significant marten population (Slauson et al. 2003), and because habitat conditions in the parks are not currently optimal for marten.

The Parks also may not be sufficiently connected to currently occupied habitat to provide for marten dispersal (Slauson et al. 2003). Even though habitat in the parks is not vulnerable to logging, the marten may be vulnerable there to other threats such as recreational disturbance and vehicle collisions. EPIC (2010) estimates that 10 percent of the California range of the Humboldt marten is on land managed by Redwood National and State Parks. (*Ibid.*).

In sum, there are no existing regulatory mechanisms at the federal level which are adequate to provide for the long-term survival of the Humboldt marten.

### ***Non-federal Lands***

Logging operations on non-federal lands in California are governed by the California Forest Practice Act of 1973 (California Public Resources Code 4511 et seq.), and the associated California Forest Practice Rules ("CFPRs") (California Code of Regulations, Title 14, 895 et seq.).

There are no existing regulatory mechanisms that protect the Humboldt marten's habitat on non-federal lands in California. EPIC (2010) estimates that approximately one-third of the California range of the marten is owned by Green Diamond Resource Company and is managed as industrial timberland. Of the private land in the Slauson et al. (2007) Humboldt marten study area, 83 percent has been logged, primarily by clearcutting.

Martens are "faring worse" on non-federal lands than on federal lands (Zielinski et al. 2001, p. 488). Slauson et al. (2007) detected martens at only 2 of 36 (5.5%) sample units on private timberlands, whereas martens were detected at 24 of 123 (19.5%) sample units on lands administered by the Forest Service where less than 15 percent of the area has been logged.

There are currently no regulations contained in the CFPRs that adequately protect the Humboldt marten or its habitat. Title 14, California Code of Regulations, 919.16 requires that non-federal landowners provide CAL FIRE with stand structure information for late successional forest stands as defined (*Ref:* 14 CCR 895.1) in order to allow the Department to assess potentially significant adverse impacts per CEQA. However, such information is only required if the proposed harvest will "significantly reduce the amount and distribution of late successional forest stands or their functional wildlife value so that it constitutes a significant adverse impact on the environment as defined in Section 895.1 [of the CFPRs]." (Title 14, California Code of Regulations 919.16(a) [939.16(a), 959.16(a)]). What's more, there are no specified protective or mitigation measures to offset any potentially significant adverse impacts to late successional forest stands contained in the CFPRs. Rather, the CFPR process for assessing impacts to late successional forest stands is simply a paper exercise that does not result in any on-the-ground protection for these critical forest types.

Finally, at this time, there are no state or federally-approved plans that would protect the Humboldt marten or its habitat on non-federal lands such as Habitat Conservation Plans, Native Communities Conservation Plans or Safe Harbor Agreements. The primary non-federal landowner in the range of the Humboldt marten, Green Diamond Resource Company, has a Habitat Conservation Plan and an associated Incidental Take Permit to cover its

activities relative to the Northern Spotted Owl (NSO) (*Strix occidentalis caurina*). This HCP does not specifically provide protections for late successional forest habitats being utilized by the marten. Thus, the conservation strategy built into the Green Diamond Resource Company HCP for the NSO provides little, if any, protection for the Humboldt marten or its habitat on the property.

In sum, there are no specified regulatory protections contained in the CFPRs for non-federal lands that would specifically address the needs of the Humboldt marten. The existing regulatory mechanisms on non-federal lands are woefully inadequate, and are a primary reason demonstrating why the Humboldt marten warrants listing under CESA.

### ***Tribal Lands***

EPIC (2010) estimates that approximately 9 percent of the California range of the marten lies within the boundaries of the Yurok reservation, and less than one percent within the Hoopa reservation. (EPIC and CBD 2010, p. 32). While the Yurok Reservation lies almost entirely within the marten's range, most of the reservation is in non-Tribal ownership, including a significant portion owned by Green Diamond. There are no publicly available data on the status of the marten on Tribal lands, thus making it difficult to assess the efficacy of what, if any, protective or regulatory mechanisms are being applied there.

### ***Summary of Existing Regulatory Mechanisms***

In conclusion, there are no existing regulatory mechanisms which adequately protect the Humboldt marten's habitat on federal or non-federal lands. The lack of meaningful regulations or other conservation measures further demonstrates that the Humboldt marten is under threat and therefore warrants listing under CESA.

### **(b) Toxicant Exposure**

Toxicant exposure is an emerging and significant threat to the survival and conservation of the Humboldt marten. Although there have been no direct studies of the potential effects of toxicant exposure on the Humboldt marten, information extrapolated from research on the effects of toxic exposure on other forest-dwelling animals demonstrates that such exposure is likely a threat to the marten.

Toxic chemicals are utilized in both domestic and agricultural settings and are usually intended to suppress populations of rodents, insects, mollusks, and other agricultural and urban pests, but can have inadvertent negative impacts on humans, pets, and other non-target animals (Erickson and Urban 2004, Albert et al. 2010, Mnif et al. 2011, Gabriel et al. 2012). Widespread secondary exposure to pesticides has been reported for raptors, carnivores, and other wildlife that consume poisoned rodents around farms and human dwellings (Albert et al. 2010, Murray 2013). Researchers have generally assumed that pesticides pose little threat to wildlife outside of agricultural and urban areas (Gabriel et al. 2013). However, a recent publication reported that 79% of fishers tested in two study areas on federal and tribal forest lands in California had been exposed to anticoagulant rodenticides (ARs), including four that died from lethal toxicosis (Gabriel et al. 2012; note: at least two more fishers in California died from AR poisoning following publication of this study: Gabriel et al. 2013). Most fishers in the study had been exposed to multiple AR compounds (range = 1–4, mean = 1.6).

ARs detected in fishers in northwestern California include brodifacoum, bromadiolone, chlorophacinone, diphacinone, and warfarin (Gabriel et al. 2012). Brodifacoum and bromadiolone are classified as second-generation anticoagulant rodenticides (SGARs). SGARs were introduced in the 1970s due to widespread development of resistance among rodents to first-generation anticoagulant rodenticides (FGARs), such as warfarin, chlorophacinone, and diphacinone (Buckle et al. 1994). ARs have also been detected in a dead NSO recovered in Mendocino County (Calforests 2014) and 34 of 84 (40%) barred owls tested for exposure in Humboldt County (Gabriel et al. 2014).

Strong circumstantial evidence implicates pervasive illegal outdoor marijuana cultivation as the primary source of pesticide exposure for forest predators in California (Gabriel et al. 2012, 2013, Thompson et al. 2014).

Large quantities of ARs, particularly SGARs, are often spread across large areas in and around illegal outdoor marijuana grow sites (Gabriel et al. 2012, Thompson et al. 2014). Gabriel et al. (2012) noted that thousands of pounds of pesticides were found at illegal outdoor marijuana grow sites in California in 2008 and that 150 pounds of pesticide were found during a single three-week eradication operation on the Mendocino National Forest in 2011. Three sites raided in Humboldt County in 2013 contained a total of at least 17 pounds of SGAR bait, which researchers estimated was sufficient to kill 2,753 woodrats, 14 fishers, or five spotted owls (Humboldt County Sheriff's Office press release). Other pesticides, such as organochlorine, organophosphate, and carbamate insecticides, some of which are banned in the U.S., are also frequently found at illegal outdoor marijuana grow sites (HSVTC 2012, Thompson et al. 2014). Pesticides are often applied along with large quantities of fertilizer at the base of marijuana plants grown outdoors (Thompson et al. 2014), suggesting that marijuana and surrounding plants may be taking up pesticidal compounds from the soil. If this occurs, then rodents and insects may accumulate pesticides through consumption of plants as well as pesticidal bait. Investigation of pathways of pesticide exposure for the marten, as well as levels of exposure and potential physiological, behavioral, and population impacts, is needed.

In summary, the potential impacts of toxic exposure on the precariously small Humboldt marten population in California is a present and ever-increasing threat. The explosion of outdoor marijuana growing and the associated use of toxic chemicals for pest suppression serve as yet another indicator that listing of the Humboldt marten under CESA is warranted.

### **(c) Climate Change**

The Humboldt marten is threatened by global climate change. Currently the climate in Humboldt marten habitat is characterized by moderate temperatures, high annual precipitation, and summer fog which support dense and continuous tree and shrub cover (Slauson et al. 2007). A change in any of these parameters resulting from climate change would threaten the survival of the Humboldt marten. As described in the Species Report:

Increased temperatures and decreased rainfall projected by climate change models within the analysis area in the short-term future (approximately 40–50 years) may result in loss, degradation, or fragmentation of suitable coastal marten habitat. Potential direct impacts to suitable coastal marten habitat include conversion of suitable forest types (i.e., moist coniferous or mixed conifer-hardwood forests) to unsuitable (for coastal martens) forest types, such as hardwood forests, and loss of the mesic, shade-tolerant shrub layer required by the coastal marten. Potential indirect impacts of climate change include the creation of an open understory due to the loss of the mesic shrub

understory mentioned above, which is preferred by coastal marten predators such as gray fox and bobcat, and thus may increase predation rates. Another potential indirect impact of climate change is the effect of a warmer and drier climate on the frequency, size, and severity of future wildfires potentially resulting in the loss, degradation, or fragmentation of suitable habitat and possibly direct mortality of coastal martens when severe wildfires burn through extant population areas; especially the SCO\_EPA and CA\_EPA where fire frequency, size, and intensity are currently a stressor on coastal marten populations. As mentioned above, coastal marten populations are already small and isolated and suitable habitat is already fragmented and greatly reduced from historical levels. Therefore, further habitat loss, degradation, or fragmentation from climate change could threaten the future viability of coastal marten populations and reduce the likelihood of reestablishing connectivity between extant populations. (USFWS 2015, at p. 57).

Climate change is therefore a threat to the conservation of the Humboldt marten.

## **IX. Degree and Immediacy of Threat**

The entire Humboldt marten population is extremely low; less than 50 individuals are believed left in California and an unknown, but presumed small and declining number, remain in Oregon. Population size is the best predictor of extinction probability. It is likely that the extant populations of Humboldt marten are below the population size needed to maintain long-term population viability, especially considering each population is reproductively isolated. Populations of at least several hundred reproductive individuals are needed to ensure the long-term viability of vertebrates with several thousand individuals being a desirable goal for many vertebrate species (Primack 1993, pp. 335–336). For a mammal their size, martens have low reproductive rates suggesting a slow recovery from population-level impacts (Buskirk and Ruggiero 1994, p. 16).

The precariously small size of the population of the Humboldt marten is a significant and immediate threat to the survival and conservation of the species. Small isolated populations are inherently vulnerable to extinction for the following four main reasons: (1) genetic problems due to loss of genetic variability, inbreeding, loss of heterozygosity, and genetic drift; (2) demographic fluctuations due to random variations in birth and death rates; (3) environmental fluctuations due to variation in predation, competition, disease, and food supply; and (4) natural disturbances resulting from single events that occur at irregular intervals, such as fires, storms, or droughts (Primack 1993, p. 255). The smaller a population becomes the more likely the factors described above in section VIII will individually or cumulatively reduce the population size even more and drive the population to extinction (Primack 1993, p. 274).

Slauson et al. (2009a) found that the probability of extinction for the Humboldt marten in their study area was higher than the probability of colonization, and that conservation actions to benefit the remnant population are therefore needed immediately. Conservation actions based solely on measures to maintain current habitat conditions for the Humboldt marten will not ensure its long-term persistence (Slauson 2003, p. 71). Given the marten's extremely small population size in California, its isolation from other populations in Oregon, its declining population status, limited range, and the variety and magnitude of threats to its continued survival, it clearly warrants CESA protection. The protection provided under CESA is necessary to prevent the Humboldt marten's extinction in California. *California Forestry Assn. v. California Fish & Game Commission* (2007) 156 Cal.App.4th 1535, 1551 ("the term "range"

as used in [CESA] refers to a species' California range”).

## **X. Impact of Existing Management Efforts**

There are currently no species-specific protective measures or management plans for the Humboldt marten. The lack of adequate protections and appropriate management planning is a primary reason why the Humboldt marten is under threat, and therefore warrants listing under CESA. Information on species and land management activities that are impacting populations of the Humboldt marten, including land classifications and uses within the range of the marten are described in the Inadequacy of Regulatory Mechanisms section, as are protective measures being taken.

Current research on the Humboldt marten is ongoing through the multi-stakeholder Humboldt marten Conservation Group, which includes independent researchers, the U.S. Fish and Wildlife Service, the Forest Service, Tribes, and Green Diamond Resource Company. To date, there is not any formal report, or other document that has been made publically available as a result of the Conservation Group’s activities.

## **XI. Suggestions for Future Management**

Hamlin et al. (2010) collected and synthesized the primary components of a conservation and management strategy for the Humboldt marten:

- Maintain all currently occupied habitat.
- Restore habitat to increase and reconnect suitable habitat patches in the vicinity of the known population (Slauson and Zielinski 2004, p. 63).
- Increase the overall size of suitable patches toward the mean size of 447 ac (181 ha) (Slauson et al. 2007, p. 466).
- Restore functional landscape connectivity to enable recolonization of suitable, but currently unoccupied habitat (Slauson and Zielinski 2003, p. 13) and establish connectivity with habitat corridors between populations.
- Establish high priority restoration areas that enlarge small suitable patches, such as late-successional conifer-dominated stands and serpentine stands with dense shrub cover, so that they exceed the minimum patch size occupied by martens [greater than 205 ac (83 ha)]. This will reconnect suitable patches currently separated by unsuitable habitat.
- Restore or maintain dense, productive shrub layers and reduce road densities in the short-term and accelerate development of late-successional stand conditions, such as large diameter live trees, multilayered canopy, and large snags and logs over the long-term (Slauson et al. 2007, p. 466).
- Develop specific stand recommendations to manage early-seral conifer stands with lower tree densities to encourage maintenance of a productive shrub layer and increase tree growth rates (Slauson 2003, p. 71).
- Protect currently suitable resting and denning structures and plan for the future recruitment of new structures (Slauson and Zielinski 2009, p. 43).
- Establish additional populations within the historical range.



## **XII. Availability and Sources of Information**

Literature cited in this petition is listed below. A disk with many of the critical documents cited will be send via U.S. Mail to the Commission along with a paper copy of the petition.

## Literature Cited

- Albert, C.A., L.K. Wilson, P. Mineau, S. Trudeau, and J.E. Elliott. 2010. Anticoagulant rodenticides in three owl species from western Canada, 1988-2003. *Archives of Environmental Contamination and Toxicology* 58(2):451-459.
- Archibald, W. R., and R. H. Jessup. 1984. Population dynamics of the pine marten (*Martes americana*) in the Yukon Territory. Pages 81-97 *In* R. Olsen, F. Geddes, and R. Hastings (editors), *Northern Ecology and Resource Management*. University of Alberta Press, Edmonton, Alberta.
- Banci, V. 1989. A fisher management strategy for British Columbia. Victoria, BC: British Columbia Ministry of Environment, Wildlife Branch. Wildlife Bulletin B-63. 117 pp.
- Blaxter, K.L. 1971. The comparative biology of lactation. *In*: Falconer, I.R., ed. *Lactation*. London.
- Bissonette, J.A., and S.S. Sherburne. 1993. Habitat preferences of unexploited pine marten (*Martes americana*) populations in Yellowstone National Park. Final report. Utah Cooperative Fish and Wildlife Research Unit, Utah State University, Logan, Utah.
- Bissonette, J. A., D. J. Harrison, C. D. Hargis, and T. G. Chapin. 1997. The influence of spatial scale and scale-sensitive properties on habitat selection by American marten. Pages 368–385 *in* J. A. Bissonette, editor. *Wildlife and landscape ecology*. Springer- Verlag, New York, New York, USA.
- Bolsinger, C. L. and K. L. Waddell. 1993. Area of old-growth forests in California, Oregon, and Washington. U. S. Department of Agriculture, Forest Service. Research Bulletin. PNW-RB-197. 29 p.
- Brown, R.N., M.W. Gabriel, S. Matthews, J.M. Higley, G. Wengart, and J.E. Foley. 2006. Pathogens associated with Pacific fishers (*Martes pennanti*) in northwestern California - final report. United State Fish and Wildlife Service, Yreka, California, USA.
- Buckle A.P., C.V. Prescott, and K.J. Ward. 1994. Resistance to the first and second generation anticoagulant rodenticides – a new perspective. *In* *Proceedings of the sixteenth vertebrate pest conference*.
- Bull, E.L., T.W. Heater, and J.F. Shepherd. 2005. Habitat Selection by the American Marten in Northeastern Oregon. *Northwest Science* 79(1): 37-43.
- Buskirk, S.W. 1994. An introduction to the genus *Martes*. *In*: Buskirk, S.W.; Harestad, A.S.; Raphael, M.G., comps. eds. *Martens, sables, and fishers: biology and conservation*. Ithaca, NY. Cornell University Press: 1-10.
- Buskirk, S.W. and L.L. McDonald. 1989. Analysis of variability in home-range size of the American marten. *Journal of Wildlife Management*. 53:997–1004.
- Buskirk, S.W. and L. F. Ruggiero. 1994. The American marten. *In*: Ruggiero, L.F., K.B. Aubry, S.W. Buskirk, L.J. Lyon, and W.J. Zielinski, (eds.). *American marten, fisher, lynx, and wolverine in the western United States*. General Technical Report, RM-254.

Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station.

Calforests (California Forestry Association). 2014. Northern Spotted Owl Science Compendium. Unpublished report submitted to the California Department of Fish and Wildlife and California Fish and Game Commission.

Calder, W.A., III. 1984. Size, function, and life history. Cambridge, MA: Harvard University Press: 431 p.

Chapin, T.G., D.J. Harrison, and D.D. Katnik. 1998. Influence of landscape pattern on habitat use by American marten in an industrial forest. *Conservation Biology* 12(6):1327-1337.

Chapin, T. G. 1995. Influence of landscape pattern and forest type on use of habitat by marten in Maine. M.S. thesis. University of Maine, Orono. 100 p.

Dawson, N. and J.A. Cook. 2009. Phylogeography of two martens (*Martes americana* and *Martes caurina*) in North America: tracking diversification in forest-associated mustelids. Abstract in 5th International *Martes* Symposium Biology and conservation of Martens, Sables, and Fishers: A new synthesis. University of Washington, Seattle, WA. 8–12 September 2009.

Douglas, R.B., and M.R. Holley. 2009. Mesocarnivore distribution on private timberlands in Mendocino County. 2008 Annual Wildlife Report. Mendocino Redwood Company, LLC.

Drew, G.S. 1995. Winter habitat selection by American marten (*Martes americana*) in Newfoundland: Why old growth? Dissertation, Utah State University. Logan, UT, 83 p.

Enders, R.K. and O.P. Pearson. 1943. Shortening gestation by inducing early implantation with increased light in the marten. *American Fur Breeder*. (Jan.): 18.

Environmental Protection Information Center, Center for Biological Diversity. 2010. Petition to List the Humboldt Marten under the Endangered Species Act.

Environmental Protection Information Center. 2010. A spatial analysis of current Humboldt marten range in California. Arcata, CA.

Erickson, W. and D. Urban. 2004. Potential risks of nine rodenticides to birds and nontarget mammals: A comparative approach. Office of Pesticides Programs, Environmental Fate and Effects Division, United States Environmental Protection Agency, Washington, D.C.

Fredrickson, R.J. 1990. The effects of disease, prey fluctuation, and clearcutting on American marten in Newfoundland, Canada. M.S. thesis. Utah State University, Logan. 76 p.

Gabriel, M.W., L.W. Woods, R. Poppenga, R.A. Sweitzer, C. Thompson, S.M. Matthews, J.M. Higley, S.M. Keller, K. Purcell, R.H. Barrett, G.M. Wengert, B.N. Sacks, and D.L. Clifford. 2012. Anticoagulant rodenticides on our public and community lands: Spatial distribution of exposure and poisoning of a rare forest carnivore. *PloS ONE* 7(7):e40163.

Gabriel, M.W., G.M. Wengert, J.M. Higley, S. Krogan, W. Sargent, and D.L. Clifford. 2013. Silent Forests? Rodenticides on illegal marijuana crops harm wildlife. *The Wildlife Society*

News. Available at: < <http://news.wildlife.org/twp/2013-spring/silent-forests/>>

Gilbert, J. H., J. L. Wright, D. J. Lauten, and J. R. Probst. 1997. Den and rest-site characteristics of American marten and fisher in Northern Wisconsin. In G. Proulx, H. N. Bryant, and P. M. Woodard (editors), *Martes: taxonomy, ecology, techniques, and management*, Provincial Museum of Alberta, Edmonton, Canada. Pp. 135-145.

Green, G.A., L.A. Campbell, and D.C. MacFarlane. 2008. Submitted. A conservation assessment for fishers (*Martes pennanti*) in the Sierra Nevada of California. USDA Forest Service, Pacific Southwest Region, Vallejo, California, 72 pages.

Grinnell, J., J.S. Dixon, and J.M. Linsdale. 1937. Furbearing mammals of California. Vol. 1. Berkeley, CA: University of California Press. 375 pp.

Hamlin, R., L. Roberts, G. Schmidt, K. Brubaker and R. Bosch 2010. Species assessment for the Humboldt marten (*Martes americana humboldtensis*). U.S. Fish and Wildlife Service, Arcata Fish and Wildlife Office, Arcata, California. 34 + iv pp.

Hanson, C.T. 2013. Habitat use of Pacific fishers in a heterogeneous post-fire and unburned forest landscape on the Kern Plateau, Sierra Nevada, California. *The Open Forest Science Journal*, 6, 24-30.

Hanson, C.T. 2015. Use of higher-severity fire areas by female Pacific fishers on the Kern Plateau, Sierra Nevada, California, USA. *The Wildlife Society Bulletin* (in press)

Hargis, C.D. and R.D. McCullough. 1984. Winter diet and habitat selection of marten in Yosemite National Park. *Journal of Wildlife Management*. 48:140-146.

Harris, L.D. 1984. *The Fragmented Forest: Island Biogeography Theory and the Preservation of Biotic Diversity*. University of Chicago Press, Chicago, IL.

Hauptman, T.N. 1979. Spatial and temporal distribution and feeding ecology of the pine marten. Pocatello, ID: Idaho State University. M.S. thesis. 84 p.

IERC 2008. Integral Ecology Research Center. 2008. Pathogens associated with fishers (*Martes pennanti*) and sympatric mesocarnivores in California. Final Report submitted to USFWS Yreka, California, 13 May 2008.

Katnik, D. D., D. J. Harrison, and T. P. Hodgman. 1994. Spatial relations in a harvested population of marten in Maine. *Journal of Wildlife Management* 58:600–607.

Koehler, G.M., J.A. Blakesley, and T.W. Koehler. 1990. Marten use of successional forest stages during winter in North Central Washington. 71:1-4.

Kucera, T.E. 1998. Humboldt marten, *Martes americana humboldtensis*. Pp. 140-141 In: Bolster, B.C., Ed. 1998. *Terrestrial Mammal Species of Special Concern in California*. Accessed May 27, 2010 at: <http://www.dfg.ca.gov/wildlife/nongame/ssc/docs/mammal/species/44.pdf>

Lettman, G. and D. Campbell. 1997. Timber harvesting practices on private forest land in western Oregon. Oregon Department of Forestry, Salem, USA.

- Lewis, J. C. and W. J. Zielinski. 1996. Historical harvest and incidental capture of fishers in California. *Northwest Science* 70:291-297.
- Lofroth, E. C. 1993. Scale dependent analyses of habitat selection by marten in the subboreal spruce biogeoclimatic zone, British Columbia. M.S. Thesis, Simon Fraser University, Victoria, British Columbia.
- Markley, M.H. and C.F. Bassett. 1942. Habits of captive marten. *American Midland Naturalist* 28:604-616.
- Mead, R.A. 1994. Reproduction in martens and fishers. In: Buskirk, S.W.; Harestad, A.S.; Raphael, M.G., comps. eds. *Martens, sables, and fishers: biology and conservation*. Ithaca, NY. Cornell University Press: 404-422.
- Millar, J.S. 1977. Adaptive features of mammalian reproduction. *Evolution*. 31: 370-386.
- Mnif, W., A.I.H. Hassine, A. Bouaziz, A. Bartegi, O. Thomas, and B. Roig. 2011. Effect of endocrine disruptor pesticides: A review. *Environmental Research and Public Health* 8:2265-2303.
- Phillips, D. M. 1994. Social and spatial characteristics and dispersal of marten in a forest preserve and industrial forest. M.S. thesis. University of Maine, Orono. 112 p.
- Potvin, F., L. Belanger, and K. Lowell. 1999. Marten habitat selection in a clearcut boreal landscape. *Cons. Bio.* 14: 844-857.
- Primack, R.B. 1993. *Essentials of Conservation Biology*. Sinauer Associates Inc., Sunderland, Massachusetts.
- Raphael, M. G., and L. L. C. Jones. 1997. Characteristics of resting and denning sites of American marten in central Oregon and western Washington. Pages 146-165 *In* G. Proulx, H. N. Bryant, and P. M. Woodard (editors), *Martes: Taxonomy, Ecology, Techniques, and Management*. Provincial Museum of Alberta, Edmonton, Alberta.
- Six Rivers National Forest. 2010. About the Forest. Accessed June 29, 2010 at: <http://fs.usda.gov/wps/portal/>
- Slauson, K.M. and W. Holden. 2009. News Release: American marten discovered in Prairie Creek Redwoods State Park: first in recent times. USDA Forest Service Pacific Southwest Research Station, California. August 25, 2009.
- Slauson, K.M. and W.J. Zielinski. 2009a. Characteristics of summer and fall diurnal resting habitat used by American martens in coastal northwestern California. *Northwest Science* 83(1):35-45.
- Slauson, K.M., J.A. Baldwin, and W.J. Zielinski. 2009b. Status and Estimated Size of the Only Remnant Population of the Humboldt Subspecies of the American marten (*Martes americana humboldtensis*) in Northwestern California. November 25, 2009. Final Report.
- Slauson, K.M., W.J. Zielinski, and K.D. Stone. 2009c. Characterizing the molecular variation among American marten (*Martes americana*) subspecies from Oregon and California. *Conservation Genetics* 10(5):1337-1341.

- Slauson, K.M., and W.J. Zielinski. 2007a. Strategic Surveys for Martes Populations in Northwestern California: Mendocino National Forest July – September 2006 Final Report. U.S.D.A. Forest Service, Pacific Southwest Research Station, Redwood Sciences Laboratory, Arcata, California.
- Slauson, K.M. and W.J. Zielinski. 2007b. Diet of American martens in Coastal Northwestern California. In: Transaction of the Western Section of the Wildlife society, Sacramento, CA. February 1-4, 2007.
- Slauson, K.M., W.J. Zielinski, and J.P. Hayes. 2007c. Habitat Selection by American Martens in Coastal California. *Journal of Wildlife Management* 71(2):458–468.
- Slauson, K. M. and W.J. Zielinski. 2004. Conservation status of American martens and fishers in the Klamath-Siskiyou bioregion. 25. Arcata, California, USA: USDA Forest Service, Pacific Southwest Research Station. Accessed June 18, 2010 at: [http://www.fs.fed.us/psw/publications/slauson/psw\\_2004\\_slauson\\_001.pdf](http://www.fs.fed.us/psw/publications/slauson/psw_2004_slauson_001.pdf)
- Slauson, K.M. 2003. Habitat Selection by American Martens (*Martes americana*) in Coastal Northwestern California. Oregon State University. Master's Thesis.
- Slauson, K.M., W.J. Zielinski, and G.W. Holm. 2003. Distribution and Habitat Associations of Humboldt marten (*Martes americana humboldtensis*) and Pacific fisher (*Martes pennanti pacifica*) in Redwood National and State Parks. Final Report. 18 March 2003. Redwood Sciences Lab, Pacific Southwest Research Station USDA Forest Service. Arcata, CA.
- Slauson, K.M., W.J. Zielinski, and J.P. Hayes. 2002. Ecology of American Martens in Coastal Northwestern California. Progress Report II. USDA Forest Service, Pacific Southwest Research Station. Redwood Sciences Laboratory, Arcata, California.
- Slauson, K., B. Zielinski, and C. Carroll. 2001. Hidden in the Shrubs: Rediscovery of the Humboldt Marten? Mountains and Rivers A Quarterly Journal of Natural History for the Klamath-Siskiyou Region. 1(2):1-12.
- Snyder, J. E., and J. A. Bissonette. 1987. Marten use of clear-cuttings and residual forest stands in western Newfoundland. *Canadian Journal of Zoology* 65:169-174.
- Spencer, W.D., R.H. Barrett, and W.J. Zielinski. 1983. Marten habitat preferences in the northern Sierra Nevada. *Journal of Wildlife Management* 47:1181-1186.
- Strickland, M.A., C.W. Douglas, M. Novak, et al. 1982. Marten. Pages 599-612 In: Chapman, J.A. and G.A. Feldhamer, eds. Wild mammals of North America: biology, management, economics. Baltimore, MD: Johns Hopkins University Press.
- Thompson, I.D. and P.W. Colgan. 1987. Numerical responses of martens to a food shortage in northcentral Ontario. *Journal of Wildlife Management*. 51: 824-835.
- Thompson, I.D. and A.S. Harestad. 1994. Effects of logging on American martens, and models for habitat management. Pages 355-367 In S. W. Buskirk, A. S. Harestad, M. G. Raphael, and R. A. Powell (editors), Martens, Sables, and Fishers: Biology and Conservation. Cornell University Press, Ithaca, New York.

Thompson, C., R. Sweitzer, M. Gabriel, K. Purcell, R. Barrett, and R. Poppenga. 2014. Impacts of rodenticide and insecticide toxicants from marijuana cultivation sites on fisher survival rates in the Sierra National Forest, California. *Conservation Letters* 7(2):91-102.

Thornburg, D. A., R. F. Noss, D. P. Angelides, C. M. Olson, F. Euphrat, and H. W. Welsh. 2000. Managing redwoods. In: R. F. Noss (ed.). *The Redwood Forest: History, ecology, and conservation of the Coast Redwoods*. Island Press, Covelo, CA. 339 pp.

U.S. Department of Agriculture; U.S. Department of the Interior [USDA and USDI]. 1994a. Final supplemental environmental impact statement on management of habitat for late-successional and old-growth forest related species within the range of the northern spotted owl [Northwest Forest Plan]. Portland, OR. 2 vol.

U.S. Department of Agriculture; U.S. Department of the Interior [USDA and USDI]. 1994b. Record of decision on management of habitat for late-successional and old growth forest related species within the range of the northern spotted owl [Northwest Forest Plan]. Portland, OR.

U.S. Department of Agriculture (USDA). 1992. Final Environmental Impact Statement (FEIS) on management of the northern spotted owl in the national forests. States of Washington, Oregon, and California. Portland, Oregon.

U.S. Fish and Wildlife Service 2015. Coastal Oregon and Northern California Populations of the Pacific Marten (*Martes caurina*): Species Report. April 2015.

Wilbert, C.J. 1992. Spatial scale and seasonality of habitat selection by martens in southeastern Wyoming. Laramie, WY University of Wyoming. M.S. thesis. 91 p.

Wynne, K. M., and J. A. Sherburne. 1984. Summer home range use by adult marten in northwestern Maine. *Canadian Journal of Zoology* 62:941-943.

Yeager, L. E. 1950. Implications of some harvest and habitat factors on pine marten management. *Transactions of North American Wildlife Conference* 15:319-334.

Zielinski, W.J., K.M. Slauson, C.R. Carroll, C.J. Kent, and D.G. Kudrna. 2001. Status of American martens in coastal forests of the Pacific states. *Journal of Mammalogy* 82(2):478–490.

Zielinski, W.J., and R. T. Golightly. 1996. The status of marten in redwoods: is the Humboldt marten extinct? Pp. 115–119 in *Conference on Coast Redwood Forest Ecology and Management* (J. LeBlanc, ed.). Humboldt State University, Arcata, California.

### XIII. Detailed Distribution Map




Figure 8.3. Coastal marten extant population areas. Central coastal Oregon (CCO\_EPA; top), southern coastal Oregon (SCO\_EPA; middle), and northern coastal California (CA\_EPA; bottom).